THE EARTH'S HYDROSPHERE

Distribution of Water on Earth

The Hydrosphere and the Hydrologic Cycle

Earth's Cryosphere


THE EARTH'S HYDROSPHERE

- The Earth's liquid water constitutes the hydrosphere.
- The vast majority of Earth's water is in the oceans (salt water), with smaller, but geologically important, quantities of fresh water in lakes, rivers, and ground water.
- The components of the hydrosphere, as well as the cryosphere (frozen water), the atmosphere, and the biosphere, participate in the global hydrologic cycle.
- Earth's water supply has had, since Earth was created, major influences on Earth's climate, its landscape and mineralogy, the composition of its atmosphere, and on the origin and evolution of life.
 - The total mass of Earth's water is about 300 times the mass of the atmosphere.
 - Without water, which facilitates the formation of carbonate rock, the atmospheric content of CO₂ would be far higher than it is.

THE EARTH'S HYDROSPHERE: Distribution of Water on Earth

	Volume	Percent of Total		
OCEANS	1,350 x 10 ¹⁵ m ³	97.3		
CRYOSPHERE (Glaciers & Polar Ice)	29 x 10 ¹⁵ m ³	2.1		
UNDERGROUND (Aquifers)	$8.4 \times 10^{15} \text{m}^3$	0.6		
LAKES & RIVERS	$0.2 \times 10^{15} \text{m}^3$	0.01		
ATMOSPHERE	$0.013 \times 10^{15} \text{m}^3$	0.001		
BIOSPHERE	$0.0006 \times 10^{15} \text{m}^3$	4 x 10 ⁻⁵		


Distribution of Water on Earth




Copyright @ 2002 by Tasa Graphic Arts, Inc.


(% of total hydrosphere)

The Hydrologic Cycle







Typical Elevation Profile of Oceanic Margins



Major Topographic Divisions and Profile of the North Atlantic Ocean Basin



The World's Ocean Floors



Copyright © 1998 Tasa Graphic Arts, Inc. All rights reserved.


THE EARTH'S HYDROSPHERE

- The hydrosphere, along with the atmosphere and cryosphere, are primarily responsible for weathering and erosion of land surfaces.
- Rain water, in combination with atmospheric CO₂, is primarily responsible for chemical weathering by carbonic acid, H₂CO₃.
- The amount of CO₂ dissolved in the oceans is much larger than that currently in the atmosphere. Since the solubility of CO₂ in water decreases with temperature, global warming could produce a positive feedback effect by releasing oceanic CO₂.
- Man-made and volcanic pollution can increase weathering by providing much stronger acids ("acid rain"; e.g. H₂SO₄), and by increasing atmospheric CO₂.
- Rain, plus the river and stream components of the hydrosphere, also provide mechanical erosion of rocks and convert them to soils and sediments.

Table 16-1 Average Surface Temperature (°C) of the Oceans Between Parallels of Latitude


North latitude	Atlantic Ocean	Indian Ocean	Pacific Ocean	South latitude	Atlantic Ocean	Indian Ocean	Pacific Ocean
70°-60°	5.60			70°–60°	- 1.30	- 1.50	- 1.30
60°-50°	8.66		5.74	60°-50°	1.76	1.63	5.00
50°-40°	13.16		9.99	50°-40°	8.68	8.67	11.16
40°-30°	20.40		18.62	40°-30°	16.90	17.00	16.98
30°–20°	¹ 24.16	26.14	23.38	30°-20°	21.20	22.53	21.53
20°-10°	25.81	27.23	26.42	20°-10°	23.16	25.85	25.11
10°- 0°	26.66	27.88	27.20	10°- 0°	25.18	27.41	26.01

Hurricanes, Typhoons and Cyclones



Global Ocean Circulation

The exchange of water, heat, and momentum between the atmosphere and ocean is responsible for the circulation in both. This leads to distinct patterns of the ocean circulation related to the distribution of winds




The ocean currents of the world with the approximate location of some of the major global wind systems.

THE EARTH'S CRYOSPHERE

- Earth's supply of frozen water, the cryosphere, is second only to the oceans in water content.
- The cryosphere consists mainly of the permanent ice caps of Antarctica and Greenland, with much smaller amounts in Arctic and mountain glaciers.
- Major changes in sea level can occur during times of global climate change (ice ages and global warming), due to associated changes in the water content of the cryosphere.
- During ice ages, glaciers can cover major parts of Earth's land area year-round for hundreds or thousands of years.
- The advance and retreat of glaciers can also produce major erosion and reconfiguration of the landscape.
- Ice ages and global warming can have major effects on the biosphere as well.


Permafrost in Land Areas


- Land areas in polar regions, such as Antarctica and Greenland, and the north slopes of Alaska and Siberia, have zones below their surfaces in which ground water remains frozen year-round.
- Regions in which soil water is permanently frozen constitute what is known as permafrost.



Continuous zone Discontinuous zone

Ice Cover of Greenland and Antarctica





Copyright @ 1998 Tasa Graphic Arts, Inc. All rights reserved.

Ice Age North Polar Coverage






Figure 9-21
Generalized geographic map of North America showing the maximum extent of glaciation in Pleistocene time. Arrows indicate direction of ice movement. [After U.S. Geological Survey.]

Sea Level Changes due to Ice Ages and Ice Cap Melting

